

Jack Rowley
GSM Association
London, United Kingdom
E-mail: jrowley@gsm.org


REFERENCES

- Dolan M, Rowley J. 2009. The precautionary principle in the context of mobile phone and base station radiofrequency exposures. *Environ Health Perspect* 117:1329–1332.
- ICNIRP (International Commission on Non-Ionizing Radiation Protection). 1998. Guidelines for limiting exposure to time-varying electric, magnetic, and electromagnetic fields (up to 300 GHz). *Health Phys* 74(4):494–522.
- ICNIRP (International Commission on Non-Ionizing Radiation Protection). 2009. Guidelines for limiting exposure to time-varying electric, magnetic, and electromagnetic fields (up to 300 GHz). *Health Phys* 97(3):257–258.
- International Agency for Research on Cancer. 1987. Asbestos. *IARC Monogr Eval Carcinog Risk Hum* 14(suppl 7):106–116. Available: <http://monographs.iarc.fr/ENG/Monographs/suppl7/Suppl7-20.pdf> [accessed 8 December 2009].
- Scientific Committee on Emerging and Newly Identified Health Risks. 2009. Health Effects of Exposure to EMF. Available: http://ec.europa.eu/health/ph_risk/committees/04_scenihir/docs/scenihir_022.pdf [accessed 15 September 2009].

ERRATUM

The authors report an error in Figure 2 of “Air Pollution Exposures and Circulating Biomarkers of Effect in a Susceptible Population: Clues to Potential Causal Component Mixtures and Mechanisms” [Environ Health Perspect 117:1232–1238 (2009)]. In “Materials and Methods,” they noted that “four influential high outliers for IL-6 > 10 pg/mL were reset to 10 pg/mL”; however, the analysis shown in Figure 2A for interleukin (IL)-6 included these outliers without them having been reset to 10 pg/mL. In all other figures showing results for IL-6 and in Supplemental Material (doi:10.1289/ehp.0800194.S1 via <http://dx.doi.org/>), the outliers were correctly reset to 10 pg/mL. The IL-6 outliers are themselves valid measurements, and the results in both cases do not change the finding that the association of IL-6 with PM_{0.25} (particulate matter < 0.25 µm in aerodynamic diameter) is stronger than associations with larger particle size fractions.

Choose from over
100 science and interdisciplinary lessons based on selected articles published by *EHP*, the top monthly journal in public, environmental, and occupational health.



International Society for Environmental Epidemiology

BEST ENVIRONMENTAL EPIDEMIOLOGY PAPER AWARD 2009

An ISEE annual award for the best environmental epidemiology paper published in a peer-reviewed journal

Award criteria include:

- **Quality (data, analysis, presentation)**
 - **Originality**
 - **Importance/impact**
- Authors, editors, and readers may make nominations. Only outstanding papers should be nominated.
 - The nominated paper should be printed in hard copy in a scientific journal between January 1 and December 31, 2009.
 - Submit nomination letters no later than **April 1, 2010** by postal mail or email, accompanied by a copy of the paper and supporting letter (maximum of 2 pages) stating why the paper should receive the award.

Address for sending nominations:

Secretariat, International Society for Environmental Epidemiology
c/o JSI Research and Training Institute, 44 Farnsworth Street, Boston, MA 02210 USA
ATTENTION: Chair, Best Environmental Epidemiology Paper Award Committee
iseepi@jsi.com

For more information visit www.iseepi.org